Volume 4: Number 2: Supplement 4: December 2020

AJTES

ALBANIAN JOURNAL OF TRAUMA AND EMERGENCY SURGERY



Focus on "Trauma and Emergency Surgery" and not only...

Official publication of

Albanian Society for Trauma and Emergency Surgery
ASTES

ISSN: 2521-8778 (print version)
ISSN: 2616-4922 (electronic version)

ACTES 2020

4th Annual Albanian Congress of Trauma & Emergency Surgery

VIRTUAL EDITION

21 – 22 December 2020, Tirana, Albania.

"Clear Vision, Careful Steps...Sure Success"

UNIVERSITETI I MJEKESISE, TIRANE

Albanian Society for Trauma and Emergency Surgery ASTES

Organized by:

UMT

(University of Medicine of Tirana) Albania.

&

ASTES

(Albanian Society for Trauma and Emergency Surgery)

Main Topics Highlights of ACTES 2020:

Politrauma
Trauma systems & Prehospital Care
Trauma Scoring system
Trauma Mass Casualties Incidents
Blunt & Penetrate Abdominal Trauma.
Chest Injuries
Tramatic Brain Injuries
Critical Care in Trauma and Emergency Surgery
Bleeding, Massive transfusion and coagulopathy
Obstetric Trauma
Pediatric Trauma
Intensive Care in Trauma
Elderly Trauma and Critical Care
Skeletal Trauma & Bone Infections
Spinal Injuries
Disaster and Military Surgery
Emergency Surgery & Soft Tissue Infections
Evolution in Emergency Medical Practices
Education in Trauma and Emergency Surgery
Sports Trauma
Surgical Critical Care & Burns

Abstract Template
Click here

Abstract Submission Is open
Click here

Abstract Submission deadline is; Monday, December 14, 2020

Congress Secretary

astes.office@gmail.com; secretary@astes.org.al; abstract.submision@astes.org.al;
Mob; +355675648310; +355696481100; +355698180015;

Volume 4; Number 2: Supplement 4: December 2020

Focus on "Trauma and Emergency Surgery" and not only...

Official publication of

ISSN: 2521-8778 (print version)
ISSN: 2616-4922 (electronic version)

M&B PUBLICATIONS



Albanian Journal of Trauma and Emergency Surgery - AJTES
Official Publication of the
Albanian Society for Trauma and Emergency Surgery - ASTES

Chairman of the editorial board
Asc.Prof. Dr. Agron DOGJANI, MD, Ph.D., FACS, FISS
University Hospital of Trauma, Tirana, ALBANIA

Editor in Chief

Prof. Dr. Kastriot HAXHIREXHA, MD, Ph.D.
State University of Tetova, Republic of North MACEDONIA

Deputy Editor

Prof. Lutfi ZYLBEARI, MD, Ph.D.
State University of Tetova, Republic of North MACEDONIA
Asc. Prof.Dr. Majlinda NACO, MD, Ph.D.
University Hospital Centre "Mother Teresa", Tirana, ALBANIA
Edvin SELMANI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA
Alfred IBRAHIMI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA
Hysni BENDO, MD.
University of Medicine of Tirana, ALBANIA

International Editorial Board

Prof. Dr.Fausto CATENA, MD, Ph.D., FACS.
Department of Emergency and General Surgery, Parma University Hospital, ITALY
Prof. Dr.Juan A. ASENSIO, MD, FACS, FCCM, FRCS *(England),* ***FISS, KM.*** - Creighton University Medical Center Omaha NE 68124 USA
Andrew BAKER, MD, Ph.D., FACS.
Orthopedic Surgeon Specialist, Durban, SOUTH AFRICA
Joakim JORGENSEN, MD, Ph.D., FACS.
University Hospital, Oslo, NORWAY
Prof. Dr.Selman URANUES, MD, FACS.
Medical University of Graz, Graz, AUSTRIA
Prof. Dr.Dietrich DOLL, MD, Ph.D., FACS.
St. Marien hospital Vechta,GERMANY
John E. FRANCIS, MD, Ph.D., FACS.
St Elizabeth Regional Health, Lafayette, Indiana, USA
Thomas B. WHITTLE, MD, Ph.D.
Vascular Surgery Specialist in Lincoln, Nebraska, USA
Prof. Dr. Sadi BEXHETI, MD, Ph.D.
State University of Tetova, Republic of NORTH MACEDONIA
Hayato KURIHARA, MD, FEBS, FACS.
Milan, ITALY
Asc. Prof. Dr.Pantelis VASSILIU, MD, Ph.D., FACS.
Faculty of Medicine, University of Athens, Athens, GREECE
Prof. Dr. Massimo SARTELI, MD, Ph.D., FACS,
Chieti Clinical Hospital "S.S.Annunziata", ASL. Lanciano-Vasto-Chieti, ITALY
Prof. Dr. Jordan SAVESKI, MD, Ph.D.
Acibadem Sistina Hospital, Republic of NORTH MACEDONIA
Carlos MESQUITA, MD, Ph.D.
Coimbra Central and University Hospital Centre, PORTUGAL
Mauro ZAGO, MD, FEBS, FACS,
Policlinico San Pietro, Ponte San Pietro, BG, ITALY
Prof. Dr.Boris SAKAKUSHEV, MD, Ph.D.
"Saint George" University Hospital, Plovdiv, BULGARIA
Prof. Dr.Bellal A. JOSEPH MD, FACS
University of Arizona, College of Medicine - Tucson, USA
Ass. Prof.Dr.Basri LENJANI MD, Ph.D.
University Clinical Center of Kosovo, Pristina, KOSOVO
Prof. Dr.Orhan ALIMOĞLU MD, Ph.D.
Faculty of Medicine, Istanbul Medeniyet University, TURKEY
Prof. Dr.Mehmet ERYILMAZ MD, MBAH
Gülhane Military Medical Academy, Ankara, TURKEY
Prof. Dr.Kenan KARAVDIC MD, Ph.D.
Clinical Center University of Sarajevo, BOSNIA AND HERZEGOVINA
Asc. Prof. Dr.Bogdan DIACONESKU MD, Ph.D.
Carol Davila University of Medicine and Pharmacy, Bucharest, ROMANIA
Ilir HASANI MD, Ph.D.
PHI UC TOARICEC Skopje, Republic of NORTH MACEDONIA
Asc. Prof. Dr.Nehat BAFTIU MD, Ph.D.
University Clinical Center of Kosovo, Pristina, KOSOVO
Asc.Prof. Dr. Nuhi ARSLANI MD, Ph.D.
UniversityMedical Centre Maribor, SLOVENIA
Gjergji SEMINI MD, Ph.D.
Ioannina, GREECE
Ass. Prof. Dr. Faton T. HOXHA, MD, Ph.D.
University Clinical Center of Kosovo, Pristina, KOSOVO

National Editorial Board

Prof. Dr. Arben GJATA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof.Dr. Xheladin DRAÇINI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof.Dr. Nikollaq KAÇANI MD, Ph.D
University of Medicine of Tirana, ALBANIA

Prof.Dr. Etmont ÇELIKU, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof.Dr. Arben BEQIRI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof.Dr. Ilia MAZNIKU, MD, Ph.D.
University "Alexander Xhuvani", Elbasan, ALBANIA

Prof.Dr. Arvin DIBRA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc.Prof. Dr. Sokol XHEPA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Dr. Edmond NUELLARI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc.Prof. Dr. Sokol BUBA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc.Prof. Dr. Aleksander HOXHA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Dr. Elizana PETRELA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Dr. Edmond PISTULLI MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Dr. Edmond ZAIMI MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Dr. Ridvan ALIMEHMETI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Dr. Rudin DOMI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Dr. Besim BOCI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc.Prof. Dr. Fadil GRADICA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Artid LAME, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Arsen SEFERI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Sokol ISARAJ, MD, Ph.D.
University Hospital Centre "Mother Teresa", Tirana, ALBANIA

Dritan TODHE, MD, Ph.D.
University Hospital of Trauma, Tirana, ALBANIA

Gëzim GALIQI, MD, Ph.D.
Shkodra Regional Hospital, ALBANIA

Leart BERDICA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Myzafer KAÇI, MD, Ph.D.
University Hospital Centre "Mother Teresa", Tirana, ALBANIA

Henri KOLANI, MD, Ph.D.
University Hospital Centre "Mother Teresa", Tirana, ALBANIA

Saimir KUÇI, MD, Ph.D.
University Hospital Centre "Mother Teresa", Tirana, ALBANIA

Shkëlzen OSMANAJ, MD, Ph.D.
University Hospital of Trauma, Tirana, ALBANIA

Zamir DEMIRAJ, MD, Ph.D.
University Hospital of Trauma, Tirana, ALBANIA

Advisory Expert

Ilir ALIMEHMETI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Assistant Editors

Gentian ZIKAJ, MD.
University Hospital Centre "Mother Teresa", Tirana, ALBANIA.

Besmira ZAMA, MD
University of Medicine of Tirana, ALBANIA.

Sindi SHANDRO
Faculty of Medicine, University of Medicine of Tirana, ALBANIA.

Seimir LAQJA
Faculty of Medicine, University of Medicine of Tirana, ALBANIA.

Mail Address: Str. Lord Bajron, 1026, Tirana, Albania
Electronic Address: E-mail: ajtesjournal@gmail.com; ajtes.editor@gmail.com

Design & Layout: Bujar KAROSHI; Web developer; Olger MELICA
Printed: M&B BOTIME & GRAPHIC STUDIO
ISSN: 2521-8778 (print version) ISSN: 2616-4922 (electronic version)

Albanian Journal of Trauma and Emergency Surgery is the official publication of Albanian Society for Trauma and Emergency Surgery - ASTES

Structure and format

AJTES is an open access, peer-reviewed periodical journal that includes editorials, reviews, original articles, casereports, hort reports, ideas and opinions, book reviews, perspectives, seminars, symposium and minisymposium, ethics and rights, health care policy and management, practice guides. The structure of each edition of the publcation comprises section categories determined by Editor and reflects the views of the Editorial Board.

Editorial expertise

Guidance from Editorial Board consists of leading authorities from a variety and respective fields of medical science. Peer review accomplished by expert selected for their experience and knowledge of a specific topic.

Important notice for authors

All manuscript must be submitted to the Editorial Office online at journal.astes.org.al/submission. Check the last page of each edition"Instruction for Authors": The work shall not be published elsewhere in any language without the written consent of AJTES. The articles published in this journal are protected by copyright, which cover translation rights and the exclusive right to repro- duce and distribute all of the articles printed in the journal. No material published in the journal may be stored on micro-films or video-cassettes or in electronic data bases and the like or reproduced photographically without the prior written permission of AJTES.

Permissions

For information on how to request permissions to reproduce articles/information's from this journal, please contact Editorial Office.

Advertising policies

The journal accepts displayed and classified advertising. Frequency discounts and special position are available. The journal reserves the right to reject any advertisement considered unsuitable according to the set policies of the journal.
The appearance of advertising or product information in the various sections in the journal does not constitute an endorsement or approval by thejournal and/or its publisher of the quality or value of the said productor of claims made for it by its manufacturer.

Disclaimer

The information and opinions presented in the Journal reflects the views of the author and not of the Journal or its Editorial Board or the Pub-lisher.
Publication does not constitute endorsement by the journal.

Official Publication of the Albanian Society of Trauma and Emergency Surgery - ASTES

Aims and Scope

Our aim is to promote interest, knowledge and quality of care in emergency and trauma surgery. ASTES was formed in 2017, it seeks to promote best practice in the provision of emergency and trauma surgery and acute care surgery, from pre-hospital care through diagnosis, intervention and intensive care to rehabilitation. This is supported by Countrywide & International collaboration, scientific research, development and delivery of training courses, and the work of the specialist sections (Polytrauma, Visceral & Chest Trauma, Skeletal Trauma & Sports Medicine, Neurosurgical, Anesthesia - Reanimation, Acute Care Surgery, ENT & Ophthalmology & Maxillofacial, Radiology, Nurse service, Disaster & Military Surgery...etc.) ASTES holds an annual scientific meeting – the Albanian Conference for Trauma and Emergency Surgery (ACTES) and produces a bi-annual journal – the Albanian Journal for Trauma and Emergency Surgery

Contents:

ORAL PRESENTATIONS *7*

INDEX *47*

ORAL PRESENTATIONS

OP - 01

COVID-19 Epidemiology in Albania during 2020

Ilir Alimehmeti

Clinical Epidemiologist & Endocrinologist

Deputy Dean – Faculty of Medicine, UMT, Tirana, ALBANIA

Abstract:

Introduction: It can be said that COVID-19 pandemic has inflicted a really severe toll in terms of infection, morbidity and mortality worldwide without sparing Albania. Moreover, during 2020 different non pharma-cological interventions where put to action in different countries and with different timing. Thus, a more clear depiction is urgently warranted. Albania presented its first confirmed cases on March 8th, which have increased to over fifty thousand until mid-December 2020, a number that is condemned to further increase reaching probably sixty thousand by the end of the year. However, such figures are only the confirmed cases registered by the Institute of Public Health that, accordingly to data from national and international sero prevalence studies, is just a frac-tion of the real number of cases during this period. Nevertheless, epidemiology is interested not only in transmission patterns and trajectories, but also on several outcome data, that in a decreasing importance order include but are not limited to mortality, inten-sive care unit hospitalization, hospital occupancy rate, morbidity, test positivity percentage etcetera. Moreover, several non-pharmacological interventions will be discussed focusing especially on their effec-tiveness in terms of incidence, based on the target population they are directed towards. ***In conclusion,*** a more clear, transparent and mean-ingful presentation of the official data will offer the participants specific easy-to-miss epidemiological insights on COVID-19 during the whole first year of the pandemic in Albania.

Keywords: COVID-19, Intervention, Transmission, Epidemiological

OP - 02

The impact of COVID-19 in Trauma and Orthopedic Surgery in Albania

Edvin Selmani

Lecturer in Orthopedic Surgery & Traumatology at the Faculty of Medicine, UMT, Tirana, ALBANIA

Orthopedic Surgeon at the University of Hospital of Trauma, Tirana, ALBANIA

Abstract

Introduction: The coronavirus 2019 global pandemic has had a significant impact on trauma and orthopaedic (T&O) departments worldwide. To manage the peak of the epidemic, orthopaedic staff were redeployed to frontline medical care; these roles included managing minor injury units, forming a “proning” team, and assisting in the intensive care unit (ICU).

Outpatient clinics were restructured to facilitate virtual consulta-tions, elective procedures were cancelled, and inpatient hospital admissions minimized to reduce nosocomial COVID-19 infections.We can say that orthopaedic training has been impacted during this period. This article discusses the impact of COVID-19 on T&O in Albania and highlights key lessons learned that may par help to proactively prepare for the next global pandemic.

Conclusions: The COVID-19 pandemic has had an unprecendented impact on T&O in Albania , with changes in outpatient clinics, fracture clinics, inpatient hospital admissions, elective and emergency opera-tive procedures, staff redeployment and reallocation of resources. The main lessons learned for the next pandemic are that orthopaedic departments need to remain flexible to infra-structural reorganisation to increase critical

care capacity, maintain a low threshold for remote staff computer access to reduce hospital staff exposure, facilitate outpatient management via telephone consultations to reduce hospital footfall, and redirect training via web-based platforms while operative experience cannot be gained.
Keywords: Coronavirus, Trauma, Global Pandemic, Orthopaedic, Training,

OP - 3

Case Report: Pneumomediastinum due to Covid -19 Pneumonia.

Fadil Gradica[1], Daniela Xhemalaj[1], Lutfi Lisha[1], Alma Cani[1], Fahri Kokiçi[1], Dhimitraq Argjiri[1], Ilir Peposhi[1], Flora Gradica[2], Valbona Rexha [2], Cleon Gradica[2]

[1]University hospital "Shefqet Ndroqi", Tirana, ALBANIA.

[2] Pharmacy

Abstract:

Introduction: Spontaneous pneumomediastinum (SP) unrelated to assisted ventilation is a newly recognised complication of severe acute respiratory syndrome (SARS). The objective of the present study was to evaluate the importance of covid -19 pneumonia, in avoiding Spontaneous Pneumomediastinum.
Case report: A 57-year- old man presented to the emergency room with 3 days of increasing cough and shortness of breath, both at rest and with minimal exertion, associated with decreased appetite and also anosmia. He denied fevers, chills, chest pains, nausea or vomiting. He did suspect recent coronavirus exposure. His chest X-ray was remarkable for bilateral patchy infiltrates. Discussion: One possible explanation of occurring Pneumomediastinum related to viral infection is that alveolar rupture may occur in SARS as a result of severe diffuse alveolar damage and cause interstitial emphysema. Air may then dissect along the broncho-vascular sheaths into the mediastinum to manifest as SP initially,which may further progr-ess to pneumotho-races and subcutaneous emphyse-ma in some patients. Pneumothorax and pneumo-mediastinum represent rare complications of Covid -19 pneumonia and only few cases of spontaneous pneumomediastinum have been published to our knowledge.
In conclusion, the spontaneous pneumomediastinum appeared to be a frequent complication of severe acute respiratory syndrome, including as well covid-19 pneumonia.
Keywords: Pneumomediastinum, Covid-19, Covid-19 Pulmonary Complication, CT-Covid Complications.

OP - 04

Abdominal Disturbances in Children with Covid-19

Ferizat Dika-Haxhirexha, Sevdije Koxha, Ledia Qatipi, Shqiponja Turkeshi, Aulona Haxhirexha

PHO "Alba- Med" – Diber, Regional Hospital – Diber, NORTH MACEDONIA

Abstract:
Introduction: In this presentation, we will present the case of three children of ages 4 to 14 years, with signs of acute abdomen. In all three patients, anamnestic data showed positive family members infected with COVID-19.
The laboratory analyzes in all of them resulted in increased values of both erythrocyte sediment and CRP, while only in one of them the values of LDH and transaminases (AST, ALT) were also increased.
Ultrasonographic examination of the abdomen in two children showed the presence of a small amount of free fluid in the abdomen and mesentery thickening with several packets of enlarged lymph nodes. Radiography of the lungs did not show pulmonary lesions in any of the children. Body temperature ranged from 37.2 to 38.50C, while urine was within the normal range.All children were hospitalized and kept under obser-vation for several consecutive days. Furthermore, at the same time they were treated with antibiotics of the group of cephalosporins of the third generation, as well as with antipyretics (paracetamol).

[illegible]

OP - 05

Management of the SARS-Cov2 Pandemic. Recommendations for the Primary Care Physician.

Hysni Bendo

General Physician [illegible]

Abstract:

Introduction: The scientific knowledge about the SARS-CoV-2 virus is increasing rapidly. The infec-tion process is very dynamic and requires permanent adjustments to the strategies for fighting pandemics at the medical and political level. The SARS-CoV2 pandemic demands a considerably high degree of activity and flexibility from general practitioners to manage their patients in the best possible way as well as to make a contribution to the protection of society and to avoid overburdening the entire health care system. The aim of this short presentation is to give recom-mendations for different scenarios and treatment occa-sions related to the SARS-CoV-2 pandemic, which are relevant for primary care physicians.

Keywords: SARS-CoV2 pandemic, Actvity, Primary care, management.

OP - 06

Nurses and Prevention of COVID-19 Infections

Aferdita Ademi, Blerim Fejzuli

Abstract:

Introduction: The purpose of this study is to manifest the extra commitments and responsibilities that nurses carry in the surgical wards to prevent outbreaks of infections with the virus COVID – 19.

Material and Methods: Patients with surgical diseases and infected with COVID - 19 from the outbreak of the pandemic until November 1, 2020, hospitalized in the Surgery Clinic of the Clinical Hospital of Tetova are the focus of the study.

Results: Out of the total number of patients hospitalized in our ward for various surgical conditions, seven of them were diagnosed with COVID-19 virus infection a few days after hospitalization. Out of total four of these patients, three of them were operated on for acute appendicitis and one with an incarcerated hernia. The infection was diagnosed in the postop-erative period, while in the other three patients, the disease was diagnosed on the second and third day after hospitalization.

Conclusion: Nurses have shown to represent a really essential part of the medical staff in dealing with the COVID-19 virus. Despite the utmost care and strict observance of all norms for preventing the spread of the virus, they too often became the victims of these infections. However, thanks to their commitment and care showed for patients, the rate of spread of infection with COVID-19 in hospital wards stayed minimal.

Keywords: Nurse, COVID-19, Prevention

OP - 07

The Role of Physiotherapy in Patients with COVID-19

Erisa Mane [1], Elda Latollari [2], Luisana Memushaj [3]

[1]FzioLife Clinic, physiotherapist, UMT, FSHMT, pedagogue, Tirana, ALBANIA

[2]Fan Noli University, lecturer, Korçë, ALBANIA

[3]FzioLife Clinic, physiotherapist, Reald University, lecturer, Vlorë, ALBANIA

Abstract:

Introduction: .

OP - 11

An Overview of the Treatment of Crush Syndrome

Merita Rroji

Department of Nephrology, University Hospital Center" Mother Tereza", Tirana, ALBANIA.

Abstract:

Introduction: Natural mass disasters affect huge popu-lations, who need basic infrastructural help and also assistance to survive. The first description of the crush syndrome appeared in the modern medical literature after the Messina earthquake in 1909. Trauma-related crush syndrome was first identified as a single/broad pathophysiologic entity in 1941. During the bombing of London, Bywaters and Beall described in detail the clinical picture in four crush cases. Three of them were oliguric and all produced dark brownish urine. All four patients died; histopathological examination of the kidneys revealed pigmented casts, polymorpho-nuclear invasion, and acute tubular necrosis (ATN). This presentation summarizes the etiopathogenesis of traumatic rhabdomyolysis and crush syndrome-based acute kidney injury. The clinical and labora-tory features, prophylaxis, and treatment of crush cases are described as well. The importance of early and dynamic fluid resuscitation is indicated for the prophylaxis of acute kidney injury. The treatment of both acute and chronic kidney diseases (CKDs) is especially problematic after disasters, because of the reason they almost always require complex technology and equipment, whereas specific drugs may be difficult to obtain for the treatment of chronic kidney patients. Although crush syndrome is a major cause of mortality in the rescued victims of massive earthquakes, the number of deaths due to crush syndrome (or fatalities of renal disaster) can be decreased by appropriate management.

Keywords: Crush syndrome, Trauma, Chronic Kidney Disease, Acute Tubular Necrosis (ATN), Fatalities.

OP- 12

Spectrum of Diaphragmatic Injury in Trauma and Emergency Radiology

Zeyad Jalee, A. A. Al-Taie, Adham Darweesh

Clinical Imaging Department, Hamad Medical Corporation, Doha, QATAR

Abstract:

Introduction:

Diaphragmatic injury (DI) is a relatively rare injury.

Mechanism of injury is either related to blunt or penetrating trauma. A75% of rupture is from blunt abdominothoracic trauma of severe grade (mainly motor vehicle accidents) and 25% from penetrating trauma (from gunshot and stab injury) 5.

BDR occurs with approximately the same frequency and injury pattern in adults and children alike

The left hemi diaphragm is three times more involved than the right, possibly due to the buffering effect of liver. The most common site of rupture is the posterolateral aspect of the hemidiaphragm between the lumbar and intercostal muscle slips.

The most commonly-her-niated viscera are the stomach and colon. Mechanisms of rupture of diaphragm include (a) sudden increase in intrathoracic/intra-abdominal pressure against a fixed diaphragm, (b) shearing stress on a stretched diaphragm, and (c) avulsion of the diaphragm from its point of attachment.

Traumatic diaphragmatic hernias are classified into three different types, depending upon the time interval for diagnosis.

Type 1 hernia - Immediately following trauma

Type 2 hernia - Within the recovery period

Type 3 hernia - When the patient presents with compli-cations related to the DI.

Complications of DI inlcude:

Gastrointensinal strangulation

Gastric volvulus

Tension gastrothorax Compression of the viscera at the site of herniation

Keywords: Diaphragmatic Injury, Hernia, Trauma, Complications.

OP - 13

Simple and Complex Pneumothorax- Diagnostic and Treatment Challenges (10 Year Study in The Regional Hospital of Vlora)

Petraq Mustaqe [1], Lorenc Gjoka [2]

[1] Surgeon in the Regional Hospital of Vlora, ALBANIA

[2] Anesthesiologist in the Regional Hospital of Vlora, ALBANIA

Abstract:

Introduction: From 2007 to 2017, 204 cases of pneu-mothorax have been diagnosed. This number is higher than before because of the CT scan and the other tech-niques that allow us to diagnose this condition easily. Most of our patients with pneumothorax, have also shown hemothorax and thoracic injuries, both pene-trating and non-penetrating. Thoracic traumas are often present in patients with polytraumas, and often these patients show pneumothorax as a complication.

Diagnostic Challenges: Considering our clinical expe-rience we have observed that there is an important discordance between the radiological response and the clinical presentation of the pat used the pleural diagnostic punction as a diagnostic procedure, this way we performed also a thoracic draining if we found air in our syringe.

Treatment challenges: In the very last years we have performed the draining procedure in the 6th intercostal space on the medio-axillary line, which has had much better results.

Conclusions: Traumatic pneumothorax is a common surgical condition. The most important causes of the thoracic traumas are: automobility accidents, falling from heights, weapon injuries and different kinds of traumas. Pathological pneumothorax is less frequent but still present. The actual treatment is almost always surgical and it consists in the positioning of a drain.

***Keywords*:** Traumatic Pneumothorax, Polytrauma, Simple Pneumothorax, Hemothorax, Medio-axillary line, Penetrating Thoracic Injury.

OP - 14

Post Operative Delirium Following Hip Surgery

Silvana Leka, Dritan Muzha, Rexhina Sturce, Alban Greca.

Anesthesia& ICU Service, University Hospital of Trauma, Tirana, ALBANIA

Abstract:

Introduction: A review of literature about incidence, early diagnosis and clinical manifestation, treatment and prevention of POD in HIP surgery (femoral neck fracture, intertrochanteric fracture).

Post-operative delirium (POD), is a variant of the delirium that is manifested in the patients who have undergone various surgical procedures and also anes-thesia, usually peaking between one and three days after their operation. It has a very high incidence in the elderly patients after repairing femoris fracture. It can cause a significant increase in morbidity, mortality, prolonged hospitalization, and medical costs. Delirium is associated with increased days of mechanical venti-lation and ICU length of stay, increased hospital length of stay, and patients' functional decline. Furthermore, all-cause mortality increases by at least 10–20% for every 48 hours of delirium. A recent meta-analysis demonstrated that delirium during hospitalization doubles a patient's risk of post-discharge institution-alization and death, and increases risk of dementia up to 10-fold. Impressively, these poor outcomes were independent of common confounders like age, sex, and comorbidities.

Additionally, in-hospital delirium is associated with more than doubled one-year medical costs. Some of the cost and mortality implications arise from the association of delirium with other postoper-ative complications, including falls, pressure ulcers, urinary tract infection, respiratory difficulties, myocar-dial infarction, and atrial fibrillation.

Keywords: Delirium, Pressure ulcer, Surgery, Femoral Neck Fracture.

OP - 15

Incidence and Treatment of Postoperative Delirium after Cardiac Surgery

Alfred Ibrahimi, Saimir Kuci, Ervin Bejko, Stavri Llazo, Jonela Burimi, Esmerilda Bulku, Edvin Prifti.

Cardiac Surgery Service. "Mother Teresa" Hospital,Tirana, ALBANIA.

Abstract:

Introduction: Delirium after cardiac surgery is a major problem. Several predisposing and precipitating risk factors have been identified for postoperative delirium. The development of delirium following cardiac surgery is associated with worse outcome in the perioperative period.

Materials and methods: during 2019 in our institution 480 cases (aged 65±19) performed CABG. 29 of them (6,1%) developed postoperative delirium. 25 of them (5,2 %) was up to 65 years. The patients were divided in two group. First, treated only with propofol and second with propofol and haloperidol. Results: mean recovery time for the first group was 3±0,5 days, and 4,5±0,9 for the second one.

Conclusion: It can be said that delirium complicates the postoperative course of a significant number of elderly patients undergoing cardiac surgery. Treatment with propofol and haloperidol has better results in recovery time.

Keywords: Delirium, Complicates, Cardiac Surgery. Propofol. Haloperidol.

OP - 16

Postoperative Complications of Coronary Artery Bypass Surgery, our Experience 2002-2020

Saimir Kuci, Alfred Ibrahimi, Ervin Bejko, Stavri Llazo, Jonela Burimi,Esmerilda Bulku, Ermal Likaj, Andi Kacani, Selman Dumani, Klodian Krakulli, Altin Veshti, Arber Aliu, Ali Refatllari, Edvin Prifti.

Cardiac Surgery Clinic at "Mother Teresa" University Hospital Center Tirana, ALBANIA

Abstract:

Background and Goal of study: To indicate the postoperative complications of bypass surgery in our clinic and to compare them with the results of the other clinics. At the same time to compare them to see how has the trend gone from 2002 to 2020.

Material and Methods: We have studied all of the patients operated with CABG, from 2002 to 2020.We have seen the percentage of myocardial infarction, low cardiac output, arrhythmias, stroke, pulmonary, renal and gastrointestinal implications are involved in the operative risk after coronary bypass surgery.

Result and Discussion: Postoperative complications after coronary bypass surgery involve all the main systems as a result of the primary pathology, accompanying disease or extracorporeal circulation effects. Complications as myocardial infarction (5-10%), low cardiac output (4-5%), arrhythmias (22-30%), stroke (3%), pulmonary (13-15%), renal (5-6%) and gastrointestinal (1,5-2%) implications are involved in the operative risk after coronary bypass surgery.

Conclusion: Myocardial function in the early postoperative period is dependent directly on the quality of surgery, type of cardio protection and operative times. Long bypass times are related to postoperative stroke and diffuse cerebral edema especially in patients with carotid stenosis.

Keywords: Arrhythmias, Stroke, CABG.

OP - 17

Management of Frailty and its Challenges in Geriatric Patients on General Anesthesia.

Majlinda Naço

Lecturer in Anesthesiology at the Faculty of Medicine, UMT, Tirana, ALBANIA.

Anesthesiologist-Intensivist at General Surgery Service at UHC Mother Theresa, Tirana, ALBANIA.

Abstract:

Introduction: In Albania there are almost 439 000 elderly people in 2021 and 1 in 25 persons performed surgery. According to deficits in function, mobility, cognition, chronic diseases and geriatric syndromes we can use clinical frailty scale, the Edmonton Frailty Scale or frailty index for calculation of frailty. We need intraoperative care, management of general anesthesia, early immobilization as well as treated of postoperative delirium because frailty increased intraoperative morbidity, increased postoperative complications especially delirium, extend hospitalization, as well as non – home discharge and mortality.

Conclusion: Many geriatric patients have multi-organ problems. Frailty is a practical, unifying concept in the care of these older people that directs attention away from organ specific diagnoses towards a more holistic viewpoint of the patient and their medical medicament.

Keywords: Frailty Syndrome, Clinical Frailty Scale, Frailty Index.

OP - 18

Computer Model of Death Probabilities in the Severe Burned Patients

Monika Belba, Majlinda Naço, Albana Aleksi, Ilir Nezha

Department of General Surgery University Hospital Centre "Mother Teresa", Tirana, ALBANIA.
Service of Anesthesiology and Intensive Care, Service of Burns and Plastic Surgery, Tirana, ALBANIA

Abstract:

Introduction: Mortality after severe burns is one of the major outcome measures in burn centres. There are many predictive factors related with the course of the disease responsible for survival after severe burn. Many burn centres have generated computer models of death probabilities based on age and TBSA (total body surface area) burned.

Purpose: To evaluate outcome of the severely burned patients treated in the burn centre and to develop a predictive model for survival from severe burns in Albania based on 1684 patients hospitalized in the Intensive Care Unit of the Service of Burns during 2009-2019.

Methods*:* The actual medical records of all acute burn patients were reviewed retrospectively. Statistical analysis was conducted using SPSS version23. We will present preliminary analysis of the logic of variables and their predictive power.

Results*:* Overall mortality in ICU is 6.8% while the mortality rate of all burns was 0.35 cases per 100000 population/year.

Conclusions: The reduction of mortality speaks for a better work of our staff to our patients. The predictive model for outcome may assist all the burn team to identify the crucial determinants of clinical outcome to establish a basis for treatment standards and to allow future comparisons of new treatment strategies. ***Keywords:*** Severe Burn, Predictive Factor, TBSA.

OP- 19

The Importance of Applying the External Clamp to Gunshot Wounds Widened

Elona HASALLA, Ilda TAKA, Blerta HASALLA, Sulejman BAHA, Elona DYBELI, Ilia MAZNIKU

University "Aleksandër Xhuvani"

OP - 20

Malignant Neuroplasies of the Uterus, Diagnostics And Imaging Methods.

Erjona ZOGAJ[2], Najada KALLASHI[1], Vidi DEMKO[1], Ina TABAKU [3], Theodor XHILLARI [2], Bukri HALILI [2]

[1]*University Hospital of Trauma, Tirana, Albania*
[2]*Radiology Technician, University Hospital of Trauma, Tirana, Albania.*
[3]*American hospital, Tirana Albania.*

Abstract:

Purpose: This topic wants to emphases the importance of medical imaging like Trans-vaginal Ultrasound, CT, PET-CT and MRI in patient with neoplasia of the Uterus. Through different technicians in early stage, we can diagnose and treat this pathology more efficiently.

Methods: This is a retrospective analytical study which is based on two study cases. Patient involved in this study was take detailed anamnesis using the standard protocol regarding the age of menopause, TZH, IMT, hypertension reproducing status and the use of actual anticoagulant

Conclusion: Uterine carcinoma is the most common malignant neoplasm of the female genital tract, the fourth most common cancer ranking after breast cancer, lung cancer and bowel cancer. Occurs mainly in post-menopausal women with an increasing incidence with increasing age.

Generally almost 2-3% of women develop malignant neoplasms of the uterus during life (life expectancy 80 years). A number of factors have recently influenced the increase in cases diagnosed with uterine cancer. These factors are related to the decrease in the inci-dence of uterine cancer in developed countries, the earlier diagnosis of cancer, the increase in life expec-tancy or the use of post-menopausal TZH.

Keywords: Carcinoma, Uterus, neoplasm, female genital tract , Imaging

OP - 21

Cranial Trauma, Pediatrics vs. Adults

.

Dorian META

Coordinator of Programs Professional in Sport and Health UMB, Tirane, ALBANIA

Abstract:

Introduction: The patient problems which have been misinterpreted as a reason for a non-good management of the information collected is common, especially in the field when it comes to trauma. One of major problems related to skull trauma (especially compared to pediatric ones) with adults is cerebral hemorrhage, where thanks to the speed of identification of the problem and with the use of the "Glasgow SCALE" and observation of well-trained students and staff. First aid can be performed in real-time from assessment by determining a fast diagnosis. Rapid Diagnosis and treatment is essential for optimal results as well as maintenance the integrity and life of the patient. ***Purpose of the study:*** Early identification of skull trauma also in children and adults and real-time definition of the traumatic event and its consequences, make up more and more a challenge as great as is necessary for the patient's life. The first line of the life chain, it is precisely the efficient and professional response in traumatic events.

Conclusions: Health personnel are required to possess specific knowledge and degrees of trauma assessment, as correct diagnosis as well as high level of knowledge affect directly in the patient's life. They have to be skilled and trained to manage hemodynamic problems and resuscitation techniques either in the use of drugs, as well as in resuscitation manipulations such as the intubation procedure.

Keywords: Skull Trauma, First Aid, Hemorrhage.

OP - 22

Adult Duplex Kidneys: Differential Diagnosis in Patients with Megaureter.

Elona DYBELI [1], Xhensila HOXHA [2], Deniona NUNCI [2], Elona HASALLA [1], Ilda TAKA [1]

[1] *Faculty of Technical Medical Sciences, Medical, "Aleksander Xhuvani" University, Elbasan, ALBANIA*

[2] *Radiology Emergency Service, University Hospital Center "Mother Teresa", Tirana, ALBANIA*

Abstract:

Introduction: Duplex collecting systems are seen in 0.7% of the healthy adult population and 2-4% of patients investigated for urinary tract symptoms. Most duplicated systems asymptomatic and diagnosed incidentally.

Case Report: We report the case of a young woman, who presented to the emergency department with right loin pain, high temperature and incontinence. Investigation by a CT scan and laboratory data revealed for a tubular formation in the right kidney, following the right ureter, and according to the clinical examina-tion was concluded for a mega-ureter and treated for two weeks with Buscopan and Ampicilin. After one month patient re-refered with the same symptoms and after that the patient underwent an open surgery. Prior to incision the patient underwent cystoscopy and stenting of the lower pole normal ureter. The patient was discharged home within five days. The stent in the lower pole ureter was removed endoscopically at routine follow-up.

Conclusion: Although those cases are hard to identify, when the situations has aggravated, radiology and surgery are the best ways to treat efficiently the case and consequently decrease the morbidity. When the patient is asymptomatic, the recommended treatment for duplex systems is ureterectmoia and heminephrectomia of the upper pole.

Keywords: Duplex Collector System, CT, Megaureter, Surgery.

OP - 23

Rehabilitation for the Replacement of Elbow Movements after Humeral Supracondylar Fractures

Ilda TAKA [1], Elona HASALLA [2], Albana SULA [3], Elona DYBELI [1]

[1] *Department of Medical Technical Specialties, "Aleksandër Xhuvani" University of Elbasan, Faculty of Medical Technical Sciences, Elbasan, ALBANIA*
[2] *Department of Preclinical Courses, "Aleksandër Xhuvani" University of Elbasan, Faculty of Medical Technical Sciences, Elbasan, ALBANIA*
[3] *Department of Nursing, "Aleksandër Xhuvani" University of Elbasan, Faculty of Medical Technical Sciences, Elbasan, ALBANIA*

Abstract:

Introduction: Supracondylar fractures are humeral bone fractures that are located slightly above the elbow. Fractures are associated with damage to ligaments, muscles, nerves and blood vessels.

Purpose: The aim of this study is to assess the importance of careful rehabilitation after bone fragment consolidation and the time required for recovery of elbow movements after treatment of humeral supra-condylar fractures in children.

Methodology: We conducted this study in 18 children injured with supracondylar fracture, of out of which 14 boys (77%) and 4 girls (23%). All are recent cases occurred and treated from 12 hours to 2 days after the initial trauma. Out of these fractures 11 cases (61%) are on the right side and only 7 cases (39%) are on the left side, given the fact that most cases use left. Elbow edema was assessed as mild in 8 cases (44%), moderate in 4 cases (22%) and severe in 6 cases (34%).

Conclusions: After gypsum immobilization, the rehabilitation begins after 3 weeks or when the frac-ture fixation is stable after 2 weeks. The movements after the completion of three weeks of rehabilitation have been complete, normal muscle trophy, preserved sensitivity. In all cases the results show a complete recovery of movements.

Keywords: Supracondylar Fracture, Immobilization, Pediatric Age, Complication.

OP - 24

Effectiveness of Physiotherapy in Children with Achilles Tendon Shortening

Erisa MANE [1], Luisana MEMUSHAJ [2]

[1] *FizioLife Clinic, physiotherapist; UMT, FSHMT, lecturer, Tirana, ALBANIA*

[2] *FizioLife Clinic, physiotherapist; Reald University, lecturer, Tirana, ALBANIA*

Abstract:

Introduction: Shortening of the Achilles tendon in children, or as it is otherwise known, Toe Walking, is a problem that is evidenced in the first year of life.

Purpose: The main objective of this study is to identify the role of physiotherapy in children with Achilles tendon shortening.

Methodology: This is a case study of a 4-year-old subject with a shortened Achilles tendon, exam-ined at 2 years of age and not treated with phys-iotherapy. The patient worked 5 times a week, 50 minutes each session according to the orthopedic soft tissue physio therapeutic protocol, lengthening and strengthening the muscles of the pulp and foot as well as active joint mobilizations with resistance. Muscle strength and length were measured. In the end it was worked with proprioception to adjust the arches of the foot.

Results: After three months of intensive physio-therapy it was found by the orthopedic doctor and physiotherapists that he gait was normalized and the endon was extended over 2cm. The arches of the foot are formed and the child escapes surgery and orthopedic shoes.

Conclusions: Physiotherapy in pediatric ortho-pedic problems is a necessity that gives signifi-cant positive results in a short time. The sooner a problem is caught, the more effective and efficient physical rehabilitation is.

Keywords: Achilles Tendon, Arches of the Foot, Pediatric Physiotherapy.

OP - 25

ERCP and its Role in the Treatment of CBD Stones

Arben BEQIRI

Surgeon at General Surgery Service, UHC "Mother Teresa", Tirana, ALBANIA.

Abstract:

Introduction: Where suitable technical equipment and trained medical and paramedical staff are available, the preferred treatment for common bile duct lithiasis (CBDL) is extraction following endoscopic sphincterotomy during endoscopic retrograde cholangiopancreatography (ERCP). In 2007, ERCP should no longer be used purely for diagnosis but rather for treating CBDL already diagnosed via non-invasive morphological examination. For patients with CBDL and the gallbladder in situ, the respective advantages and disadvantages of preoperative, peri-operative and postoperative endoscopic treatment should be considered case by case, according to the team's preference and the equipment available. In cases involving a surgical arrangement, which has altered gastroduodenal anatomy, a papilla that is hard to reach, large stones or associated stenosis, special endoscopic techniques must be used. Techniques for the supply and management of anticoagulants and antiplatelet drugs in collaboration with the cardiologist and anaesthetist must be fully mastered. Endoscopic biliary sphincterotomy is not without risk (acute hemorrhagic pancreatitis, haemorrhage, perforation, infection) and this must be explained to the patient and family as well as the expected benefits. If ERCP is to be used increasingly in the treatment of CBDL, it is essential that medical practitioners are trained in how to recognize and take immediate action with respect to any signs of complications.

Keywords: Lithiasis, ERCP, Risk.

OP - 26

Postoperative Complications after Appendectomy: A 10-Year Experience of the Department of Surgery of University Hospital of Patras

Françesk MULITA, Levan TCHABASHVILI, Fotios ILIOPOULOS, Nikolas DRAKOS, Ioannis KEHAGIAS

Department of Surgery, General University Hospital of Patras, GREECE

Abstract:

Introduction: Appendicitis is one of the most common abdominal conditions in both children and adults, requiring emergency surgery. Appendectomy (laparoscopic or open approach) is the gold-standard treatment for acute appendicitis. Postoperative complications such as wound infection and intra-abdominal abscess formation can be seen after appendectomy. The aim of this study was evaluated postoperative complica-tions after open appendectomy (OA) and laparoscopic appendectomy (LA).

Materials and Methods: From the time of January 2008, to December 2017, records of patients with appendicitis were retrieved from computer database of our institution for analysis. Patients' demographic data as well as perioperative and postoperative outcomes of LA and OA group were compared.

Results: During the study period, 1037 appendectomies were performed (706 LAs and 331 OAs). Wound infection was found in 0.28 % (2/706) of patients who underwent LA and 3.32 % (11/331) of patients of OA group ($p<0.05$). In addition, the postoperative intra-abdominal abscess occurred in 3.26% (23/706) of patients of LA group and 3.63% (12/331) of patients who underwent OA ($p>0.05$).

Conclusion: This retrospective study, shows that there was no measurable difference in incidence

OP - 27

Some consideration about Incarcerated Abdominal Hernia.

Dritan ÇOBANI [1], Agron DOGJANI [2], Henri KOLANI [3].

[1] *Department of General Surgery University Hospital of Trauma, Tirana Albania.*

[2] *University of Medicine of Tirana, Department of Surgery, Tirana, Albania.*

[3] *Department of General Surgery UHC Mother Theresa", Tirana, Albania.*

Abstract

Incarcerated hernias are characterized by anamnesis with local tumefaction accompanied by clinical data of pain or intestinal transit blockage, accompanied by clinical and local data confirmed by basic and advanced imaging examinations such as plain radiography, CT or sometimes abdominal ultrasonography.

Classical repair techniques are the most commonly used techniques for inguinal hernias, however recently other surgical modalities such as the use of a prosthetic mesh have become widely accepted.

Material and Methods: the patients in this study were grouped according to the type of hernia during the admission (inguinal, umbilical, incisional, femoral). The patients were evaluated in terms of risk factors affecting their health condition such as age, gender, concomitant diseases, and the presences of intestinal strangulation and necrosis.

Results: Inguinal hernia was the most frequent cause of incarcerated abdominal wall hernia especially in males, followed by umbilical, incizional and finally femoral hernia which was most frequent in females. The rate of intestinal resection due to strangulation and necrosis was found significantly higher among femoral and inguinal hernias as compared to the other types of hernia (umbilical and incizional). Concomitant disease, strangulation, advanced age (> 65 years), and time from the onset of symptoms were found to have significant influences on morbidity and mortality.

Conclusion: Incarcerated abdominal hernias are surgical emergencies with high morbidity and mortality rates. This is the reason that in the patients.

Keywords; Incarcerated hernias, intestinal transit blockage, repair techniques

OP - 28

Influence of early clinical data on the selection of appropriate tactics and surgery in the management of gastric cancer

Imri VISHI, Riza IDRIZI, Sylejman METUSHI, Ragip BYTYQI, Sami SPAHIU, Nijazi VISHI, Afrim TASHOLLI, Xheladin REQICA, Mufail HASANI

Department of Surgery, Regional Hospital, Ferizaj, KOSOVO

Abstract:

Introduction: Stomach cancer is the second most common cause of cancer deaths worldwide, and one of the most common cancers. Due to its often occult clinic, it is unfortunately usually diagnosed in the advanced stages of the disease, which is of prognostic importance in treatment outcomes.

Material and methods: This presentation takes into account the clinical data on different stages of stomach cancer, which consist of data obtained from our General Surgery Service, during the period 2016-2018…Result; The data, after being filed and recorded in a pre-prepared form, were taken from the preoperative anamnesis of 87 patients treated for stomach cancer ...

Conclusions: In this study, the most common data recorded in patients diagnosed with gastric cancer were weight loss and gastrointestinal hemorrhage (melena and hematemesis), which were directly related to an advanced stage of the disease, conditioning thus in the selected surgical technique(total gastrectomy, subtotal, with removal of attacked lymph nodes, with adjuvant and neo-adjuvant therapy) and consequently the prognosis ...

Keywords: Gastric cancer, Melena, Hematemesis, gastrectomy

OP - 29

A Surgeon's View of Abdominal Rectus Muscle Endometriosis, A Case Report and Literature Review

Faton T. Hoxha [1, 2], Rrustem Musa [2], Albi F. Hoxha [4], R. Mati [3]

[1]University of Gjakova "Fehmi Agani, Medical Faculty, Gjakova, KOSOVO

[2]University Clinical Hospital Service of Kosovo, Surgery Clinic, Prishtina, KOSOVO

[3]Institute of Anatomic Pathology, Medical Faculty, University of Prishtina, KOSOVO

[4]Trakya University, Medical Faculty, Edirne, TURKEY

Abstract:

Introduction: Abdominal rectus muscle endometriosis after gynecologic surgery is a rare pathological entity, defined as endometrial tissue superficial to the peritoneum. Often is misdia-gnosed and referred to surgeons for treatment. Case presentation: We report a Cauca-sian female, 34 years old, six years after a cesarean (Pfannenstiel incision), with abdominal discomfort, intermittent pain in lover abdomen, nausea, with palpable and sensi-tive mass above the scar last six months.

Treatment consisted in wide surgical full thickness abdominal wall resection, including the involved abdominal rectus muscle, followed defect by prosthetic abdominal wall repair. Through our case we discuss literature review.

The postoperative course was uneventful, HP revealed abdominal rectus muscle endometriosis with free margins.

Conclusions: Abdominal rectus muscle endometriosis is an unusual and unique form of endometriosis with diagnostic and treatment delay. Proper combination between clinical data, Sonography color Doppler find-ings, may substantially upgrade correct preoperative diagnosis of abdominal wall endometriomas.

Keywords: Abdominal Wall Endometriosis, Rectus Muscle Endometriosis, Color Doppler Sonography, Abdominal Patch Grafting

OP - 30

Incisional Hernias and Treatment Options

Kastriot HAXHIREXHA [1], Aulona HAXHIREXHA[2]

[1]Clinical Hospital, Tetovo, NORTH MACEDONIA

[2]Department of Pastic Surgery University Hospital Centre "Mother Theresa", Tirana, Albania.

Abstract:

Intraduction: Incisional hernias continue to be a significant problem in general surgery. The causes of incision hernias after a surgical intervention are numerous,
while the main ones are operative wound infection, obesity, male sex, diabetes, BPH, etc.
Classical repair techniques are the most commonly used techniques for incisional hernias, however recently other surgical modalities such as the use of a prosthetic mesh have become widely accepted.

Results: The actual study included 57 patients operated on from incisional hernia in the periodfrom January 2018 to January 2020.The age of the patients included in the study ranges from 27 to 73 years.
Most of the patients, respectively, 31 of them were male, while only 26 were female. In terms of BMI, nine of the patients were classified as normal weight (18.5-24.9), 23 were overweight (BMI 25.0 -29.9), 19 patients were first-degree
over-weight (BMI 30.0 - 34.9), while six patients were classified with class II obesity (BMI 35.0-39.9). Correction of the incisional hernia in 21 patients was performed by closing the defect without the use of prosthetic materials (herniorrhaphy). In 27 patients, hernioplasty was performed using polypropylene mesh, while in nine of them, the correction of the defect was performed through the composite dual mesh.
Conclusion: Correction of incisional hernias with the use of prosthetic materials marks a lower rate not only of recurrence but also of other postoper-ative complications, especially pain, and shows a higher degree of quality of life. In this context, the use of dual composite mesh shows higher safety indices in terms of hernia recurrence.

Keywords: Hernia, BMI, hernioplasty, polypropylene mesh

OP - 31

Emergency Medical Care and Management of Sports Injuries on the Football Court

Basri Lenjani [1]*, Premtim Rashiti [1]*, Gani Shabani [1], Arber Demiri [1], Besarta Pelaj [2], Erza Voca Mulaj [3], Arban Demi [3], Dardan Lenjani [3]

[1]*Emergency Clinic, University Clinical Centre of Pristina, KOSOVO*

[2]*College Medical of Sciences Resonance*

Abstract:

Introduction: Sports medicine is a clinical subspecialty that deals with the examination, monitoring, diagnosis, treatment and prevention of injuries that occur during sports events, training and physical activ-ities in pre-hospital settings.

Purpose: Providing emergency medical care at all basic stages of managing minor and multiple injuries on the football field and in other sports in order to implement BLS, ACLS, BTLS, PTLS, ATLS care measures reducing morbidity, disability and mortality.

Working methodology: The research is of retrospec-tive, descriptive, qualitative type. The material was taken from the archive of the Emergency Clinic of UCCK for the period January-December 2019.

Result: Sports injuries are very costly, and according to the pathology with diseases were 13 cases or 18.57%, injuries were 57 cases or 81.43%. The number of athletes with current illnesses or injuries in sports matches according to the age of 15-20 years was 27 cases or 38.58%, 20-25 years were 28 cases or 40.00% and 25-35 years were 15 cases or 21.42%

Discussion and conclusions: Education of medical staff, nurses, paramedics with courses, use of medical equipment, BLS, ACLS, BTLS, PTLS, ATLS and size and longer follow-up are required for plausible evidence and systematization of treatment principles.

Keywords: Calcaneal Fracture, Wajdi, Complication

OP - 32

A Rare Complication of Calcaneal Fracture – Calcaneal non-Union, Case Report

Andri Sherehii, P.P. Popyrkanich

Uzhgorod district clinical hospital of Uzhhorod district council of Zakarpattia region, UKRAINE

Abstract:

Introduction. The most vulnerable to such injuries are physically active, mostly male, able-bodied persons.

The purpose. Our research had the intention to study published reports of postoperative intra-articular calcaneus fractures, determinate the main reasons for such complications and study the ways to improve the treatment results.

Material and methods. We retrospectively studied anamnestic data, medical history from 2018, and the treatment process of our patient in 2019-2020 with calcaneus ORIF post-operative non-union at Trauma department of Uzhhorod Hospital.

Case report: A 56 years old male patient with over 20 years smoking experience received injury as a result of falling from a height in 2018. Next day he was admitted to the trauma department of the City Hospital with Diagnosis: Closed calcaneus intraarticular AO 82 C2 (Sanders3) fracture, with joint depression and fragment displacement. Applied ORIF with restoration of the posterior talar articular surface of the calcaneus and LCP fixation on 12th day after the trauma, followed by

protected weight bearing for 9 weeks.

Discussion and review of the literature. It can be said that Wajdi's study substantiates the possibility of using the Bone Marrow Concentrates or Platelet Rich Plasma methods.

Conclusions. Non-union after Calcaneus fracture ORIF management is an extremely rare complication. size and longer follow-up are required for plausible evidence and systematization of treatment principles.

Keywords: Calcaneal Fracture, Wajdi, Complication.

OP - 33

The Role of Skeletal Traction on the Acutely Reduced post Traumatic Hip Dislocation

Ervin NEPRAVISHTA [1], Tahir CELA [2], Eris NEPRAVISHTA [3]

[1, 2] *Department of Orthopaedics and Traumatology*

[1, 2] *University Hospital of Trauma, Tirana. ALBANIA*

[3] *Department of Physiology, Aleksandër Moisiu University, Durrës*

Abstract

Introduction: The condition of post traumatic hip dislocation, is a very common pathology, that is actually encountered in our emergency department. If not properly approached it may lead to very potential and sometimes even life threatening complications. Our routine protocol includes the immediate skeletal traction after successful closed reduction, to every case of post traumatic hip dislocation referred to our centre. The scope of this article is to challenge our protocol; once again emphasise the importance; and to concentrate on easing the difficulties faced during management of this condition, including proper diagnosis, classification, and the up-to-date treatment protocols. Special consideration is focused on the role of skeletal traction on avoiding long term and disabilitating complications such as the AVN of femoral head, taking into account the most updated international guide-liness and best on-the-field investigator recommendations.

Keywords: Traumatic hip, Dislocation, Treatment protocols.

OP - 34

Fragility Fractures- Current Concepts

Artan BANO

Orthopaedic Surgeon, University Hospital of Trauma, Tirana, ALBANIA

Abstract

Introduction: The treatment of the fragility fractures is a challenge for the Orthopaedic Surgeon. Poor bone quality, high degree of comminution, patient co-morbidities and need for long term rehabilitation are some of most important factors affecting morbidity and mortality rate. Despite these difficulties, new techniques and also special implants design have been designed, in order to face even the most difficult cases. Based on recent data recommendations for treatment of fragility fractures have been developed having final target 3 column strategy: early diagnosis and treatment, good rehabilitation and secondary prevention.

Keywords: Fracture, Technique, Treatment

OP - 35

External Fixation of Femoral Shaft Fractures

Anxhela AHMATAJ [1], Zamir DEMIRAJ [2]

[1] *Physiotherapist, University Hospital of Trauma, Tirana, ALBANIA*

[2] *Orthopeadic Surgeon, University Hospital of Trauma, Tirana, ALBANIA*

Abstract:

Introduction: Conversion of temporary external fixation to an intramedullary within the first two weeks after a femoral shaft fracture is standard practice. However, due to financial constraints, in large parts of the world external fixation of femoral shaft fractures is often the definitive treatment.

Out of 13 fractures, 10 of them were followed up for a minimum period of 39 weeks. The average follow-up time was 75 weeks. 7 fractures were closed, and 6 open. 9 fractures united at an average of 31 weeks.

There were two non unions, one of them infected. Secondary surgical procedures were performed for two non unions and in three cases of delayed union. One re-fracture occurred, which was successfully treated with repeat external fixation. Only 5 patients regained full range of motion. The average flexion was 72°. Pin tract infections occurred in 3 patients. Satisfactory results can be obtained with definitive external fixation of femoral shaft fractures.
Pin tract infections, although a common occurrence, are not a major problem and can be treated by local wound care and antibiotic therapy.
The most common problem is significant decrease in the range of motion of the knee.
Keywords: External Fixation, Shaft Fracture, Flexion, ROM.

OP - 36

Femoral Neck Fractures

Zamir Demiraj, llir Hasmuca, Anxhela Ahmataj

Orthopedic Services University Hospital of Trauma, Tirana, ALBANIA.

Abstract:
Introduction: Femoral neck fractures are very frequent pathologies that can be found in patients of all age groups; however, the most effected group are elderly patients. In the before-mentioned age group, these pathologies, are presented as a consequence of a low-energy trauma which, in most cases, occurs in a domestic setting. As for adults and young adults, neck of the femur fractures are presented as an outcome of a high energy trauma, such as: vehicular accidents or a high fall. When studying these fractures, a series of classifications exist, of which the most used in clin-ical practice are: Garden, Pauwels and the AO-group. These classifications have an important role in deciding the method of treatment and defining the prognosis. The vascularization of the head of the femur also is an essential factor in deciding the method of treatment and the definition of the prognosis.

The methods of treatment are overall quite famous: osteosynthesis and endoprosthesis. In elderly patients endoprosthesis is the best method of treatment, though the same cannot be said for adults and young adults.
For these age groups osteosynthesis is the recommended option, however this treatment intakes two major future risks: pseudoarthrosis and avascular necrosis. Taking into account the previously mentioned factors, a correct decision of the treatment protocol is the golden key to a successful recovery.

Keywords: Neck Fracture, Endoprosthesis, Avascular Necrosis, Osteosynthesis.

OP - 37

NSAIDs Provide Effective Analgesia in Traumatic and Orthopedic Patients, but can Cause Acute Kidney Injury, AKI.

Vrenos Hodaj [1], Grisilda Gjana [2]

[1] *Internist, Vice director of University Trauma Hospital,Tirana, ALBANIA*
[2] *Nephrologist, Internal Medicine Service/ SB2, Tirana, ALBANIA*

Abstract:
Case report: Flurbiprofen causes AKI, acute interstitial nephritis.
Introduction: We will present an elderly woman, who received three single doses of Flurbiprofen 100mg, for three days in a row for body and joint pain. A 69-year-old woman presented with fatigue, lower limb edema, purpura, oligo-anuria, and elevated PA values. Signs appeared after the patient received the Flurbiprofen. Urinalysis: leukocytes 8-10/ field, erythrocytes 4-5/ field and 6.6gr% proteinuria. Urine eosinophil screening was positive. A biopsy of the skin was taken. Empirical therapy with prednisolone 1mg/kg weight was initiated that started on the first day of hospitalization. After treatment the elements on the skin began todisappear. The creatinine level began to normalize, and the values got up to 1.3 mg/dl. Skin biopsy did not show inflammatory elements. The patient was discharged from the hospital after 14 days. After 1 month, the patient had no cutaneous elements and renal function was normal, creatinine 0.9mg/dl, urea 35mg/dl, but proteinuria was present. The Next follow- up was three months later and she was found in a good condition with normal urea and creatinine levels as well.

Results: The use of NSAID-s leads to renal damage, increases need for dialysis, length of hospital stay and increases possibility of mortality.

Keywords: AKI, NSAID-s, AIN.

OP - 38

Spontaneous intracranial hematoma - when urgent intervention is necessary?

Artur Xhumari

Lecturer in Neurosurgery at the Faculty of Medicine, UMT, Tirana, ALBANIA

Abstract:

Introduction: When a spontaneous intracerebral hematoma is disclosed in a CT scan, there are many different pathologies to be differentiated, further exams to be ordered and treatment options to be taken into consideration.

Through some latest operated cases we highlight the logical steps to undertake and discuss the pearls and pitfalls of managing intracerebral hematomas.

Metastasis, cerebral aneurysm, hypertensive hematoma, GBM are some of etiologies we will discuss. When taken into consideration the age, comorbidities, antecedents, anamnestic elements, then the imaging suggests the next steps to take for the etiological diagnosis as well as surgical or conservative management.

Keywords: Hematoma, CT Scan, Etiological

OP - 39

Role of Decompression and Neurolysis in Late Surgery for Neonatal Brachial Plexus Palsy

Ridvan Alimehmeti [1], Florian Dashi [1], Arian Çelaj [2] Arba Cecia [3], Mishel Qirinxhi [1], Gramoz Braçe [1]

[1]Department of Neuroscience, UHC"Mother Teresa", University of Medicine, Tirana, ALBANIA

[2]Department of Orthopedics, University of Medicine, UHC "Mother Teresa", Tirana, ALBANIA

[3]George Washington University, 2121 I St NW, Washington DC. 20052, USA.

Abstract:

Introduction: It is generally accepted that patients with neonatal brachial plexus palsy (NBPP) who do not recover within the first year of age should undergo microsurgery.

Methods: We treated 10 cases for neonatal brachial plexus palsy in an age interval from 4 to 17 years. The criteria for supraclavicular brachial plexus (SCBP) microsurgical decompression and neurolysis were based on the presence of positive Tinel sign over SCBP and the presence of almost complete passive movement in extra-rotation and abduction of the arm.

Results: All 10 cases of NBPP were selected with above-mentioned criteria to undergo SCBP microsurgical exploration. During surgery adherent and thick epineural layer was found to constrict the upper and middle trunk fascicular elements. All of the operated cases benefited both sensitive and motor improvement in a year follow-up.

Discussion: We believe that the fibrotic epineurium does not follow the same growth pace as the neural fascicles that are contained within. With time this incongruence in growth ends up in compression of the fascicle

Conclusions: Microsurgery for decompression and neurolysis of brachial plexus in late NBPP should be considered under good selection criteria.

Keywords: Neurolysis, Palsy, Microsurgery.

OP - 40

Corneal Topographic Changes After Subscleral Trabeculectomy

Marsida Krasniqi[1], Ahmed Abdel Kader Farrag[2], Ahmed Mahmoud Amin [3], Ahmed Ismail Mohamed Kamel [3], Mohamed Gamalattia El Ashkar [3].

[1]Department of Medicine, University Aleksander Moisiu, Durres, ALBANIA

[2] *Department of Ophthalmology, El Matarya Teaching Hospital, Egypt and International Eye Clinic, ALBANIA*
[3] *Department of Ophthalmology, Faculty of Medicine, Al Azhar, University EGYPT*

Abstract:

Introduction: Glaucoma is a cause of permanent blindness in older adults. Although glaucoma can be initially managed by conservative treatments, trabeculectomy has traditionally been considered the "gold standard" for filtration surgery.

Aim of the study: is to investigate the effect of trabeculectomy on corneal topography, anterior chamber depth, and aberrations by a SIRIUS topography.

Patients and methods: The present or actual study was a prospective cohort study including 30 eyes of patients who were scheduled to undergo trabeculectomy at Ophthalmology Department of Al-Azhar University Hospital from January 2018 to August 2019.

Results: We compared included eyes according to site of flap midway (between K1&K2) versus vertical flap (on K). We found no statistically significant differences between studied groups in terms of K1 ($p = 0.29$), K2 ($p = 0.33$), and astigmatism values ($p = 0.38$). there was statistically significant difference between pre and postoperative values of ACD ($p < 0.001$). Eyes showed statistically significant decrease in ACD postoperatively.

Conclusion: The trabeculectomy surgery can cause changes in keratometry values, aberrations and ACD. These changes can be significant enough to affect visual acuity, the accuracy of IOL power calculation, and refractive outcomes after combined or future cataract surgery.

Keywords: Glaucoma, Corneal Topographic Changes, Subscleral Trabeculectomy.

OP - 41

Perioperative Assessment in Renal Failure

Suela Mumajesi

Department of Nephrology, University Hospital Center" Mother Tereza", Tirana, ALBANIA.

Abstract:

Introduction: Chronic kidney disease is becoming an increasingly major problem for society and health care. The increase in the quality of life and longevity has made the number of people with CKD mark galloping steps. Today in America 1 in 7 people suffers from renal failure.

CKD patients during their lifetime undergo surgical interventions at least one time, so it is very important to evaluate pre- and post-operative risks. Early recognition of chronic renal disease is very important, as well as pre- and post operative risks not only in renal patients but also those at high risk of developing AKI which goes up to 1% in the healthy population. The strategy evaluation of renal failure patients and those with high risk plays an essential role in the outcome during surgical interventions. That's why numerous studies have been conducted on the assessments of pre- and post operative risks in patients with renal failure. Hence, the focus on this topic is very important in clinical practice."

Keywords: CKD, Renal Failure, Surgical Intervention

OP - 42

Advanced trauma life support training- How useful it is?

Agron Dogjani [1, 2*], Arben Gjata [1,] Kastriot Haxhirexha [3], Amarildo Blloshmi [4], Sindi Shandro [1], Seimir Laqja [1].

[1] *University of Medicine of Tirana, Albania*
[2] *University Hospital of Trauma, Tirana, Albania,*
[3] *State University of Tetovo, R of North Macedonia.*
[4] *National Centre of Medical Emergency Tirana, Albania*

Abstract;

For more than 20 years, following the changes that took place in Albania in 1992, we receive information from trauma patients in the media about the delivery of trauma services.

Police records are worthy of a war balance report. Trauma is estimated to be the cause of deaths in 13.7% / 100,000 people. Every two days three Albanian Citizens die because of Automobile Accidents…Changes have already taken place, but the question arises, who cares about the trauma and its management? Formerly and now the state ... ok ... but? Is this enough?
The trauma service delivery is now evolving along the lines of a central and spoken model with a concentration of expertise and specialization in the center surrounded by smaller units that feed from the center.
The study showed a 19% increase in the chances of survival since the introduction of these changes. Another 1,600 trauma victims are alive today due to developm-ents in the administration of trauma patients in England over the past six years.
In this system or mode of organizing what is today the NCME or organizing the transport of traumatized from the scene to the University Hospital of Trauma, by medical staff who have done a substantial part of the PhTLS® course. How is educational status of personal in other regions far from Capital? ATLS® student courses in our country have been implemented by few and individually… ASTES, by default rules, has acquired the right to organize ATLS® in Albania…

Keywords: Trauma, ATLS, Course, Skills

OP - 43

The Role of Sleeve Bronchial Resection in the Treatment of Lung Cancer

Fatmir Caushi, Ilir Skenduli, Danjela Xhemalaj, Alban Hatibi, Eljana Shima, Fari Kokiqi, Alma Cani, Rinald Kortoci, Ilir Peposhi

Department of Thoracic Surgery of University Hospital "Shefqet Ndroqi", Tirana, ALBANIA

Abstract:

Introduction: Sleeve resection was first introduced in 1947 by Prince-Thomas. Allison in 1959 reported first sleeve lobectomy with pulmonary artery construc-tion. Sleeve lobectomy was considered inferior to pneumo-nectomy, but work by Ferugson et al and Deslauriers et al showed that sleeve lobectomy has better outcome and lower morbidity and mortality as compared to pneumonectomy.

The indication for a sleeve resection for lung cancer is well established: a tumor arising at the origin of a lobar bronchus precluding simple lobectomy, but not infiltrating as far as to require pneumonectomy. Bronchial sleeve lobectomy is reported to be adequate for 5% to 8% of patients with resectable lung cancer but rates as high as 13% have been reported recently.

Indications:

Tumor invading/protruding main stem bronchus.

As an alternative to pneumonectomy in patients with poor cardiopulmonary reserve.

Endobronchial bronchogenic carcinoma.

Carcinoid tumors and also low-grade malignancy like bronchial gland carcinomas if complete resection can be obtained.

Contraindications:

Complete resection of tumor not achievable by bron-chovascular sleeve resection N2 disease

Our experience: In the last year we have performed with very good results one bronchial sleeve resection and two semi sleeve bronchial resections for lung cancer with histopathologic diagnosis squamous carcinoma.

Keywords: Sleeve Resection, Lung Cancer, Lobec-tomy, Pneumonectomy

OP - 44

Bricker Diversion after Radical Cys-to-prostatectomy. 10 Years Experience

Gëzim Galiqi [1], Shpëtim Ymeri [1], Albert Pesha [2], Shkelqim Ferko [2], Bilbil Hoxha [3], Anton Ndoj[2], Imri Vishi [4]

[1] Regional Hospital of Shkoder, ALBANIA.
[2] American Hospital of Tirana, ALBANIA.
[3] Ikeda Clinic Tirana, ALBANIA.
[4] Regional Hospital of Ferizaj, KOSOVO.

Abstract:
Introduction: Radical cyst-prostatectomy represents the standard treatment for invasive bladder cancer. Since 1950 when Bricker introduced his technique of

diversion, it has been a standard of care in most of the cases after radical cyst prostatectomy.

Objectives: Evaluating the early and long-term compli-cations of Bricker diversion in our institution for a period of 12 years with 125 patients.

Materials and Methods: We evaluated 125 patients from 2008 to 2020. There were 24 women and 91 men. We evaluated post operatory complications linked with diversion and also long term complications because the patient came on a regular follow up.

Results: In terms of early the complication the most common was ileus which resolved with liquids and nasogastric tube 4-5 days after surgery. We had 2 patients who were reoperated because of intestinal fistula. In one of them we made an ileostomy and he died the 7 day after second intervention. Most common long term complication was stricture of the anasto-mosis of ureters with Bricker ileum. We diagnosed this problem in 15 patients.

Conclusions: Bricker diversion remains an effective way after radical cystectomy for bladder invasive carcinoma in female, in advance diseases of bladder and in old patients. Such complications can be treated effectively.

Keywords: Bricker, Cysto-prostatectomy, Complications

OP - 45

Lung Cancer Immunohistochemistry Predictors.

Leart Berdica 1,2 Teona Bushati 1,2

[1] *Department of Anatomic Pathology and Forensic Medicine, Medical Faculty, UMT, Tirana, ALBANIA*
[2] *Department of Pathology, American Hospital, Tirana, ALBANIA*

Abstract:
Introduction:The identification of certain genomic alterations (EGFR, ALK, ROS1, BRAF) or immunological markers (PD-L1) in tissues or cells has led to targeted treatment for patients presenting with late stage or metastatic lung cancer.

Immuno-histochemistry is a widely available techniq-ue that is less challenging and can provide clinically meaningful results quickly and cost-efficiently in comparison with other tech-niques. In addition, immuno-histochemistry allows for the evaluation of cellular localization of proteins in the context of tumor structure. In an era of precision medicine, pathologists are required to classify lung cancer into specific subtypes and assess biomarkers relevant to molecular-targeted therapies.

This review summarizes the hot topics of immuno-histochemistry in lung cancer, including adeno-carcinoma vs squamous cell carcinoma; neuro-endocrine markers; primary lung carcinoma vs metastasis, ALK, ROS1, and EGFR; (iv) PD-L1 (CD274); (v) Major pitfalls in evaluating immuno-histochemical results are also described.

Keywords: WHO Classification, Morphology, ALK, Immuno-histochemistry, Non-small cell lung carcinoma (NSCLC), Small Cell Lung Carcinoma (SCLC), Metastasis, Predictive Biomarkers.

OP - 46

Endovenous Laser Ablation

Skënder Veliu [1], Jasmina Kröpfl [2], Dominika Vrbnjak [3]

[1]*Surgeon, Surgical ward, General Hospital dr. Jože Potrč Ptuj, SLOVENIA*
[2] *Nurse for controlling hospital infection General Hospital dr. Jože Potrč Ptuj SLOVENIA*
[3]*Assistant Professor University of Maribor Faculty of Health Sciences Maribor, SLOVENIA*

Abstract:
Introduction: The chronic venous insufficiency is a significant public health problem in the modern world. It is most commonly manifested as varicose veins. Approximately 20-25% of the population experiences some problems with varicose veins. There are many approaches with which the symptoms of the chronic venous insufficiency can be alleviated.Surgical treatment is the only method that can permanently remove the reflux in the superficial venous system.

Endovenous laser ablation (EVLA) is the most used method of treatment of varicose veins. This method utilizes the heat of a laser to ablate a vein. It is used to permanently remove the reflux in the greater and lesser saphenous vein. This way, we accomplish that there is no blood flow in the treated veins. Their function is taken over by deep veins of lower extremities. Advantages of this technique are a fast re-convalescence, a significantly lower number of complications such as inflammation or bleeding, and no need for hospitalization of the patient. This paper presents the basic characteristics of the minimally-invasive thermal ablation techniques – EVLA, the experiences and results of treatment using this method in General hospital dr. Jože Potrč since 2016.

Keywords: Chronic Venous Insufficiency, endovenous Laser Ablation, Experience.

OP - 47

Is Emergency Carotid Endarterectomy in Patients with Severe Symptomatic Carotid Stenosis Justified?

Sokol Xhepa

Service of Vascular Surgery, University Hospital Center "Mother Teresa" of Tirana, ALBANIA

Abstract:
Introduction: Carotid intervention for recently symp-tomatic, severe carotid stenosis should be regarded as an emergency procedure in patients who are neurolog-ically stable, and should ideally be performed within 48 hours of a transient ischemic attack (TIA).
Exclusion criteria or cases with a very high perioperative risk for stroke are observed in patients with: not clear time of onset of symptoms, Rankin score ≥ 3, ischemic infarction > 2/3 of middle cerebral artery territory, no neurological "plateau", fluctuation in consciousness and intracerebral haemorrhage on brain imaging.
Three "old" rules to avoid intra-operative complications are: accurate surgical technique, early heparin - ization and early clamping internal carotid artery.
The adjunctive risks in very early carotid endarterectomy are: reperfusion of brain zones with loss of auto-regulation, hyper perfusion, haemorrhagic conversion of acute infarct and increased perilesional oedema.

The benefits of the urgent carotid intervention are: removing a source of ongoing embolism, preventing the early progression toward thrombosis, improving overall cerebral perfusion and reducing cumulative neuronal loss by restoring blood flow to the ischemic penumbra.

Natural history studies suggest that 8 – 15 % of patients with 50 – 99 % stenosis will suffer a stroke within 72 hours of their index symptom (Ross Naylor). Clinical and anatomic characteristics of individual patients dictate the identification of the subgroup of patients that need emergency carotid endarterectomy and the definition of timing.
Keywords: Stenosis, Intervention, Carotid, Endarterectomy

Peer Review System

Peer Review System

AJTES conducts evaluation of work in the manuscripts by peers who have an expertise in the relevant field. A thorough peer review will be carried out with a motto to maintain standards of quality, improve performance and provide credibility. Our goal of performing a comprehensive peer review system is to publish only the research results that are suitable for a scientific journal.

AJTES implements double-blinded peer review system to ensure unbiased review.

We ensure that all the articles published in Acta Scientific undergo integrated peer review by peers and consequent revision by authors when required.

Why peer review? What are the benefits?

The author receives detailed and constructive feedback from experts in the field.

The process can alert authors to errors or gaps in literature they may have overlooked. It can assist with making the paper more applicable to the journal readership.

It may enable a discussion (between the author, reviewers, and editor) around a research field or topic.

Readers can be assured that the research they are reading has been verified by subject experts.

Peer review system of AJTES assures that

1. Submitted article is original work which has not been previously published nor isunder consideration by another journal, in part orwhole;
2. The article meets all applicable standards ofethics;
3. The paper is relevant to the journal's aims, scope, andreadership;
4. A submitted article presents original researchfindings;
5. A submitted article offers a comprehensive critical review and evaluation of keyliterature sources for a given topic;and
6. The article is methodologically and technicallysound.

Article processing flowchart

Editor in Chief Guidelines

An Editor-In-Chief plays a major role, holding the responsibility of all the Journal operations and policies. The editor in chief is held accountable for efficient management of the Journal.

Roles and Responsibilities

With a great faith and trust in the professionalism of the elected person, we expect that our Editor-In-Chief will raise the reputation and profile of the Journal with respect to its competitors in the international marketplace by fulfilling the following responsibilities.

Content Accuracy

- ◊ Ensures accuracy, fairness and completeness of all the content.
- ◊ Ensures that all the corrections and clarification policies of AJTES are carried out and are presented promptly.
- ◊ Reviews feedback/evaluations of articles and other content and take steps to avoid future mistakes and hold the articles that are not ready for publication.
- ◊ In collaboration with other Editors, decides the appropriate placement of content online.
- ◊ Co-ordinates daily editing and online printing.
- ◊ Works closely with other Editors in determining the Journal features.
- ◊ Reads and edits all the content on the web pages of AJTES.
- ◊ Writes Articles and Editorials as assigned.
- ◊ Works with Managing Editor and other Editors to develop web components.
- ◊ Has final authority in the publication matters.

Planning and Association

- ◊ Maintains a master calendar and day book file for planning of overage and follow-up of works on a timely basis.
- ◊ Meets all deadlines and other requirements expected of all the Journal members.
- ◊ Conducts weekly Editor meetings online.
- ◊ Convenes Editors as needed for planning and problem solving.
- ◊ Makes frequent evaluations to ensure the publication is meeting the needs and interests.
- ◊ Provides leadership and supervision to other Editors.
- ◊ Monitors day to day work activities.

Guidance for Maintaining Quality

- ◊ Assist in the selection of Editors and Reviewers to join the Panel as required.
- ◊ Implementations to grow the Journal.
- ◊ Determine and implement strategies for increasing quality submissions to the Journal.
- ◊ Promote the Journal and encourage high quality submissions.
- ◊ Encourage appropriate revision of manuscripts and accept only high-quality papers that are high impact and contribute new and important information to the field.

Criteria for Editor-In-Chief

- ◊ An experience with broad knowledge in the relevant field.
- ◊ Extensive networks and high standing in the research community.
- ◊ Strong publication record.
- ◊ Deep understanding of peer review.
- ◊ Decisive and fair.
- ◊ Excellent organizational skills and ability to meet deadlines.
- ◊ Demonstrated vision for AJTES and the ability to think and act strategically towards the vision.
- ◊ Highly motivated and committed to AJTES.
- ◊ Time to undertake the role effectively.

Pausing your Editorship

We understand that other commitments may impact on your ability to perform your duties. It is therefore possible to pause your participation as an Editor-In-Chief. If you would like us to temporarily refrain from contacting you with Journal-related requests, just let us know beforehand, so that we can work and arrange for an alternative to avoid disturbance in smooth functioning of the Journal.

As this role is very crucial for effective running of the Journal, the Editor-In-Chief is expected to be available as needed.

The contributions of editors will be a valuable fruit to the journal.

Cover Letter

Manuscript Title: _____

_____

I/We certify that I/we have participated and contributed sufficiently for the completion of the manuscript and have agreed to have my/our name listed as a contributor.

I/We believe that the manuscript represents valid and credible work. Neither the content of this manuscript nor any other unified content with substantially similar or comparable substance under my/our creation or authorship has been published/is constantly considered for publication elsewhere. I/We certify that complete data regarding the study is proclaimed in this manuscript and no data from the study have been/will be published solely or separately. I/we attest that, I/we will cooperate in providing any required data/information related to the study during peer review by the editor or their assignees. Any direct or indirect financial interests or conflicts that exist or may be perceived to exist have been disclosed in the cover letter. Sources of external support for this work are mentioned in the cover letter.

I/We hereby exchange all copyright proprietorship, including all rights incidental thereto, solely to AJTES, in the issue that such work is published by AJTES.

AJTES grants the rights to the corresponding author to incorporate any necessary changes and he/she will act as the guarantor or surety for the manuscript on our behalf.

All persons who have made substantial and significant contributions to this work, but are not contributors, are mentioned in the acknowledgement with the written permission of the contributors to include their name. If there is no acknowledgement part in the manuscript, that means I/we have not received any contributions and also, no contributor has been omitted.

I/We have no conflicts of interest to disclose.

If you feel that the manuscript is appropriate for your journal, I/We suggest the following reviewers:

Name	Email	Affiliation	Country

Corresponding Author details

Name:

Department:

University/Organization:

Country:

Phone Number:

Email:

List of Authors

S. No	Name	Email	Signature

I certify that the above information is correct, and I would like to submit manuscript towards the journal for publication.

Corresponding Author

(Name, Signature)

AJTES - Manuscript Template

Journal Name

Type of article (E. g: Research article, Review article, Mini-Review, Case Report, Short Communication, Opinion, Letter to Editor, Symposium, Commentary, Perspective, Conceptual Paper, Proceeding etc…)

Title of the Article

Authors (List here the names of all the authors)

Authors[1], Author[2] and Author[3]

[1]Author's Department, University, Country

[2]Author's Department, University, Country

[3]Author's Department, University, Country

***Corresponding author**

Corresponding author name, Affiliation, Address, City, State, Country, Tel; Fax; E-mail:

Email:

Author 1:

Author 2:

Albanian Journal of Trauma and Emergency Surgery
AJTES
EDITOR

1. Abstract

The abstract should be clear, relative, descriptive, self-explanatory and no longer than 400 words. Do not include references or formulae or any special character in the abstract.

2. Keywords

Provide minimum 5-6 keywords.

3. Abbreviations

If there are any abbreviations in the article, they have to be mentioned.

4. Introduction (Level 1 heading)

Introduction should reflect the background, purpose and significant of the study that is carried out.

AJTES - Manuscript Template

5. Materials and Methods

All the materials and methods that are used to complete the study should be mentioned.

6. Results and Discussion

Results and discussion must illustrate and interpret the reliable results of the study.

7. Conclusion

Conclusion should reflect and elucidate how the results correspond to the study presented and provide a concise explanation of the allegation of the findings.

8. Acknowledgements

A short acknowledgement section can be written acknowledging the sources regarding sponsorship and financial support. Acknowledging the contributions of other colleagues who are not included in the authorship of this paper should also be added in this section. If there are no acknowledgements, then this section need not be mentioned in the paper.

9. Conflict of interest

Declare if any financial interest or any conflict of interest exists.

10. Bibliography

The heading of the references section must not be numbered.

References should be in MLA format and also provide the link for all the references.

Mention the reference numbers within square brackets in the body text of the article.

E.g.: Esteban R., *et al.* "Photoprotective implications of leaf variegation in *E.dens-canis* L. and *P. officinalis* L". *Journal of Plant Physiology* 165.12 (2008): 1255-1263. http://www.ncbi.nlm.nih.gov/pubmed/18180073

AJTES - Manuscript Template

Note: If there are any sub headings in the body text, sub-categorize them accordingly under the heading in which they fall.

For example:

1. Heading

1.1. Sub-heading

1.1.1. Sub-sub-heading

Please go through the below tabular description to prepare your manuscript.

Style Name	**Short Description**			
	Font size (pt)	***Font style***	***Bold (or) Normal***	***Alignment***
Title	14	Times New Romans	Bold	Center
Author	12	Times New Romans	Bold	Justify
Affiliation	11	Times New Romans	Normal	Justify
Headings	11	Times New Romans	Bold	Left
Sub-heading	10	Times New Romans	Bold	Left
Sub-sub-heading	10	Times New Romans	Bold	Left
Body text	11	Times New Romans	Normal	Justify
Table text	10	Times New Romans	Normal	Justify
Table Legend	10	Times New Romans	Normal	Justify
Figure Legend	10	Times New Romans	Normal	Justify
Bibliography	11	Times New Romans	Normal	Justify


Manuscript Submission
Screening for Unique Content
Unique Content above 80%
Unique Content below 80%
Article Modification
Review of Article by Reviewers
Review Comments sent to Author
Revision of Article according to Review comments
Revised Article sent to Reviewers
Approval of Revised Article
Rejection of Revised Article
1st Review Stage
Review of Revised Article by Editor
Review Comments sent to Author
Revision of Article according to Review comments
Revised Article sent to Editor
Approval of Revised Article
Rejection of Revised Article
2nd Review Stage
Galley Proof
Online Publication

AUTHOR PUBLICATION FEE

Article Processing Charge (APC)

The authors should note that costs are involved in every stage of the publication process, like manuscript handling blind-review, form submission to publication, copy-editing, typesetting, tagging and indexing of articles, internet expenses electronic composition and production, web-media charges of the final article on dedicated servers, electronic archiving to ensure global dissemination and administrative and overheads, and other recurring expenses, the author is asked to pay an article publication fee. The publication fee is obligatory for online publication.

AJTES has designed the Article Processing Charge (APC) in a very nominal and affordable manner such that it will be feasible and convenient for all the researchers to publish their eminent work without any barrier or obstacle. Please go through the below list regarding the processing charge.

AJTES charges a nominal Publication fee of 200 euro (Inclusive of all taxes) after the acceptance of article for publication.

For ASTES members, we charge 150 euro after the acceptance of article for publication. Mode of Payment

	Bank Account Details
Account Name:	AJTES JOURNAL
Account Number:	902019123003047803
Bank Name:	ALPHA BANK
IBAN:	AL5690211364019123003047803O
SWIFT Code:	CRBAALTRXXX

*The article accepted for publication will be processed further towards publication after the payment of the publication fees.

Withdrawal Policy: An author is free to withdraw article at no charge within a week of its original submission. However, once peer review is completed and paper is accepted for publication, then Article will be processed for journal issue. Hence author(s) are requested to pay about 30% of the publishing fee for withdrawal of Article, as a compensation for the strenuous process involved in it. For any queries/ discussions regarding manuscript withdrawal/article processing fee, please contact us at: ajtes.editor@gmail.com



Section 1 of 2

Become an Active ASTES Member 2020

The names and email addresses entered in this form will be used exclusively for the stated purposes of the society and will not be made available for any other purpose or to any other party.

Email address *

Valid email address

This form is collecting email addresses. Change settings

ter section 1 Continue to next section

Section 2 of 2

Additional Data

Description (optional)

Name Surname *

Short-answer text

Telephone Number *

Short-answer text

Address *

Long-answer text

Institution *

Short-answer text

Profession *

Long-answer text

Title

- [ ] Professor
- [ ] Associate Professor
- [ ] Doctor of Sciences
- [ ] PhD
- [ ] Medical Doctor
- [ ] Master of Sciences
- [ ] Other...

Country *

Short-answer text

5th Annual Albanian Congress of
Trauma & Emergency Surgery

ACTES 2021 - Main Topics

Politrauma

Trauma systems & Prehospital Care

Trauma Scoring system

Trauma Mass Casualties Incidents

Blunt & Penetrate Abdominal Trauma.

Chest Injuries

Tramatic Brain Injuries

Critical Care in Trauma and Emergency

Surgery Bleeding, Massive transfusion and coagulopathy Obstetric Trauma

Pediatric Trauma

Intensive Care in Trauma

Elderly Trauma and Critical Care

Skeletal Trauma & Bone Infections

Spinal Injuries

Disaster and Military Surgery

Emergency Surgery & Soft Tissue Infections

Evolution in Emergency Medical Practices

Education in Trauma and Emergency Surgery

Sports Trauma

Surgical Critical Care & Burns



10th EDITION

ATLS® Student course

✓ 01-03 Nëntor 2021

ATLS® Instructor course

✓ 04-05 Nëntor 2021

ATLS®

Advanced Trauma Life Support®

ATLS®
Inaugurale Course
Albania

Përmbledhje e ATLS® course

**Përbëhet nga testi teorik para & pas ATLS® Course.*
**Të dhëna teorike thelbësore për traumat*
**Diskutimi rastesh interaktive në Klasa-laborator në vartësi nga skenaret.*
**Leksione & prezantime & Diskutime interaktive të rasteve me trauma të #.*
**Performancë përfundimtare & vlerësimi i aftësive teorike praktike të kursanteve*
**Pas përfundimit të kursit,*
- Pjesëmarrësit duhet të ndjehen të sigurtë në zbatimin aftësitëve të mësuara në ATLS® course

Target Grup:

** Mjekë* **Specializantë* ***Studentë të Mjekësise*

**Mjekë me experiencë klinike*

**Të gjithë profesionistët e shëndetit*

- ***VENDET e limituara...***
- ***FEE = 500* & 300**Euro***

coordinator.atls.alb@gmail.com
atlsalbania2018@gmail.com

www.astes.org.al

+355699749467; +355693465944

Albanian Society for Trauma end Emergency Surgery

THE COMMITTEE ON TRAUMA

Index Table

A

A. A. Al-Taie 11
Adham Darweesh 11
Aferdita Ademi 10
Afrim Tasholli 21
Agron Dogjani 4, 7,12, 21, 28, 46
Ahmed Abdel Kader Farrag 28
Ahmed Ismail Mohamed Kamel 28
Ahmed Mahmoud Amin 28 Albana
Aleksi 17
Albana Sula 30
Alban Greca 16
Alban Hatibi 25
Albert Pesha 25
Albi F. Hoxha 22
Alfred Ibrahimi 15
Ali Refatllari 15
Alma Cani 8, 25, 33
Altin Veshti 15
Amarildo Blloshmi 28
Andi Kacani 15
Andri Sherehii 17
Anxhela Ahmataj 20, 28
Arba Cecia 24
Arban Demi 18
Arben Beqiri 20
Arben Gjata 10, 13
Arber Aliu 15
Arian çelaj 24
Artan Bano 19
Artur Xhumari 23
Aulona Haxhirexha 7, 22
Avdyl Pacolli 12

B

Basri Lenjani 13, 18
Besarta Pelaj 18
Bilbil Hoxha 25
Blerim Fejzuli 10
Blerta Hasalla 31
Bujar Millaku 12
Bukri Halili 32

C

Cleon Gradica 8

D

Daniela Xhemalaj 8
Danjela Xhemalaj 25
Dardan Lenjani 18
Deniona Nunci 31
Dhimitraq Argjiri 8
Dominika Vrbnjak 27
Dorian Meta 29
Dritan Cobani 21, 46
Dritan Muzha 16

E

Edvin Prifti 15
Edvin Selmani 7
Elda Latollari 8
Eljana Shima 25
Elona Dybeli 30, 31
Elona Hasalla 24, 30, 31
Eqrem Gara 13
Erisa Mane 8, 29, 33
Eris Nepravishta 18
Erjona Zogaj 32
Ermal Likaj 15
Ervin Bejko 15
Ervin Nepravishta 18
Ervis Cara, 24
Erza Voca Mulaj 18
Esmerilda Bulku 15

F

Fahri Kokiçi 8
Fari Kokiqi 25
Fatmir Caushi 25
Faton T. Hoxha 22
Ferizat Dika-Haxhirexha 7
Florian Dashi 24
Fotios Iliopoulos 21
Françesk Mulita 21

G

Gani Shabani 18
Gëzim Galiqi 25
Gramoz Braçe 24
Grisilda Gjana 19

H

Haki Dragusha 12
Hysni Bendo 9

I

Ilda Taka 30, 31
Ilia Mazniku 31
Ilir Alimehmeti 9
Ilir Nezha 17
Ilir Peposhi 8, 25
Ilir Skenduli 25
Imri Vishi 22, 29, 47
Ioannis Kehagias 21
Isuf Bajrami 12

J

Jasmina Kröpfl 27
Jonela Burimi 15

K

Kastriot Haxhirexha 10, 13, 22
Klodian Krakulli 15

L

Leart Berdica 26
Ledia Qatipi 7
Levan Tchabashvili 21
llir Hasmuca 20
Lorenc Gjoka 14
Luisana Memushaj 8, 29, 33
Lutfi Lisha 8

M

Majlinda Naço 16, 17
Marsida Krasniqi 28
Matjaž Horvat 13
Merita Rroji 12
Mishel Qirinxhi 24
Mohamed Gamalattia El Ashkar 28
Monika Belba 17
Mufail Hasani 21
Muhamet Petreva 12

N

Najada Kallashi 32
Nijazi Vishi 21
Nikolas Drakos 21
Nuhi Arslani 13

P

Petraq Mustaqe 14
P.P. Popyrkanich 17
Premtim Rashiti 18

R

Ragip Bytyqi 21
Rexhina Sturce 16
Ridvan Alimehmeti 24
Rinald Kortoci 25
Riza Idrizi 21
R. Mati 22
Rrustem Musa 22

S

Saimir Kuci 15
Sami Spahiu 21
Seimir Laqja 10, 13
Selman Dumani 15
Sevdije Koxha 7
Shkelqim Ferko 25
Shpëtim Ymeri 25
Shqiponja Turkeshi 7
Silvana Leka 16
Sindi Shandro 10, 13
Skënder Veliu 27
Sokol Xhepa 27
Stavri Llazo 15
Suela Mumajesi 23
Sulejman Baha 31
Sylejman Metushi 21

T

Tahir Cela 18
Teona Bushati 26

Theodor Xhillari 32

V

Vidi Demko 32
Vrenos Hodaj 19

X

Xheladin Reqica 21
Xhensila Hoxha 31

Z

Zamir Demiraj 20, 28
Zeyad Jalee 11

Albanian Society for Trauma and Emergency Surgery

ASTES

Organised by:

Albanian Society for Trauma and Emergency Surgery (ASTES)

www.astes.org.al

5-th Annual Albanian Congress of Trauma and Emergency Surgery

November 12-13, 2021
Tirana, Albania

SAVE THE DATE!

"With persistence and clear ideas, anything is possible...